

AD

RSIC-811

NUMERICAL SOLUTION OF HYPERBOLIC EQUATIONS AND SYSTEMS WITH TWO INDEPENDENT VARIABLES BY A METHOD OF THE RUNGE-KUTTA TYPE. I

by

Nguyen Kong Tuy

Vestsi Akademii Navuk Belaruskay SSR,
Seryya Fizika Matematychnykh Navuk,
No. 4, pp. 110-120, 1965.

Translated from the Russian

June 1968

This document has been approved for public
release and sale; its distribution is unlimited.

REDSTONE SCIENTIFIC INFORMATION CENTER REDSTONE ARSENAL, ALABAMA


JOINTLY SUPPORTED BY



U.S. ARMY MISSILE COMMAND



GEORGE C. MARSHALL SPACE FLIGHT CENTER



N 68-34456
(ACCESSION NUMBER)

(THRU)

25
(PAGES)

(CODE)

TMX 61184
(NASA CR OR TMX OR AD NUMBER)

19
(CATEGORY)

FACILITY FORM 602

GPO PRICE \$ _____

CSFTI PRICE(S) \$ _____

Hard copy (HC) _____

Microfiche (MF) _____

ff 653 July 65

DISPOSITION INSTRUCTIONS

*Destroy this report when it is no longer needed.
Do not return it to the originator.*

DISCLAIMER

*The findings in this report are not to be construed as
an official Department of the Army position unless so
designated by other authorized documents.*

24 June 1968

RSIC-811

**NUMERICAL SOLUTION OF HYPERBOLIC EQUATIONS
AND SYSTEMS WITH TWO INDEPENDENT VARIABLES
BY A METHOD OF THE RUNGE-KUTTA TYPE. I**

by

Nguyen Kong Tuy
Seryya Fizika Matematychnykh Navuk,
No. 4, pp. 110-120, 1965.

Vestsi Akademii Navuk Belaruskay SSR,

Translated from the Russian

This document has been approved for public
release and sale; its distribution is unlimited.

Translation Branch
Redstone Scientific Information Center
Research and Development Directorate
U. S. Army Missile Command
Redstone Arsenal, Alabama 35809

NUMERICAL SOLUTION OF HYPERBOLIC EQUATIONS AND SYSTEMS WITH TWO INDEPENDENT VARIABLES BY A METHOD OF THE RUNGE-KUTTA TYPE. I

by

Nguyen Kong Tuy

Discussed is the application of two-iteration Runge-Kutta algorithms in the solution of the Cauchy problem for hyperbolic equations and systems with two independent variables.

The question of the application of Runge-Kutta type methods to the numerical solution of partial differential equations has been considered by several authors [1-3].

In the present article, use is being made of two-iteration Runge-Kutta algorithms in the solution of the Cauchy problem for hyperbolic equations and systems with two independent variables. The results for one equation $u_{xy} = f(x, y, u, u_x, u_y)$ with Cauchy conditions along the line segment $x - y = \text{const}$ are generalized for the case of a system of such equations and are being used for the solution of a nonlinear Monge-Ampere equation. An analogous problem with Cauchy conditions along a curve segment will be discussed later.

1. Statement of the Problem

Let us consider the equation:

$$u_{xy} = f(x, y, u, p, q) \quad , \quad (1)$$

(u is an unknown function of x and y ; $p = u_x$; $q = u_y$) with initial conditions given along the segment AB of the straight line $x + y = \text{const}$, in the form:

$$u^0 = u^0(x) \quad , \quad p^0 = p^0(x) \quad , \quad q^0 = q^0(x) \quad . \quad (2)$$

u^0, p^0, q^0, f are considered to be continuous and differentiable a sufficient number of times.

Let us construct in the region ABC of definition of the solution a uniform rectangular grid with step h (Figure 1). We will distribute nodes of the grid in layers parallel to the initial line segment AB. In making calculations on the k^{th} layer, for simplicity, we will denote the earlier found values u, p, q , in the nodes of the $(k - 1)^{\text{st}}$ layer, as well as the initial data in the problem (1), (2), as u^0, p^0, q^0 .

We will introduce the following designations: $f(x, y, u, p, q) = F(x, y)$, $F(x_1, y_1) = F(M)$.

For the elementary triangle MNP of the grid with vertices $M(x_1, y_1)$, $N(x_2 = x_1 + h, y_2 = y_1 - h)$ on the $(k - 1)^{\text{st}}$ layer and vertex $P(x_2, y_1)$ on the k^{th} layer (Figure 2) the following relationships are known:

$$u(P) = u^0(x_1) + \int_{x_1}^{x_1 + h} p^0(x) dx + \iint F(x, y) dx dy, \quad (3.a)$$

$$p(P) = p^0(x_2) + \int_{y_1 - h}^{y_1} F(x_2, y) dy, \quad (3.b)$$

$$q(P) = q^0(x_1) + \int_{x_1}^{x_1 + h} F(x, y_1) dx, \quad (3.c)$$

where the multiple integral in (3.a) is being taken over the region of MNP.

Let us take as an initial point the point M, corresponding to $h = 0$. Then, using (3) we can find an expansion in powers of h at the point M for the increments $\Delta^* u = u(P) - u(M)$, $\Delta^* p = p(P) - p(M)$, $\Delta^* q = q(P) - q(M)$ in the form

$$\begin{aligned} \Delta^* u = & hp^0(x_1) + \frac{h^2}{2!} \frac{dp^0(x_1)}{dx} + \frac{h^3}{3!} \frac{d^2 p^0(x_1)}{dx^2} + \frac{h^2}{2!} F(M) \\ & \left[+ \frac{h^3}{3!} \left[2 \frac{\partial F(M)}{\partial x} - \frac{\partial F(M)}{\partial y} \right] + O(h^4) \right], \end{aligned} \quad (4.a)$$

$$\Delta^* p = h \frac{dp^0(x_1)}{dx} + \frac{h^2}{2!} \frac{d^2 p^0(x_1)}{dx^2} + hF(M) + \frac{h^2}{2!} \left[2 \frac{\partial F(M)}{\partial x} - \frac{\partial F(M)}{\partial y} \right] + O(h^3) , \quad (4.b)$$

$$\Delta^* q = hF(M) + \frac{h^2}{2!} \frac{\partial F(M)}{\partial x} + O(H^3) . \quad (4.c)$$




FIGURE 1




FIGURE 2

In the present article the problem is being set forth of constructing Runge-Kutta type algorithms, which give approximate values of the increments Δu , Δp , Δq .

2. Formal Algorithms of the Runge-Kutta Type

Values of the function $F(x, y)$ used in the i^{th} iteration ($i = 1, 2$) for the calculation of Δu , Δp , Δq , will hereafter be denoted as F_i , F_i^+ , F_i^- , respectively. Let the values u , p , q at the point p , obtained after the i^{th} iteration, be u^i , p^i , q^i , and $\int_{x_1}^{x_1+h} p^0(x) dx = I(h)$ [for the calculation of $I(h)$ see Paragraph 4].

At the first iteration we assume that:

$$u^1 = u^0(x_1) + I(h) + \frac{1}{2} h^2 F_1 , \quad (5.a)$$

$$F_1 = \omega_1 F(M) + \tau_1 F(N) ; \quad \omega_1, \tau_1 \geq 0 , \quad \omega_1 + \tau_1 = 1 , \quad (5.b)$$

$$p^1 = p^0(x_2) + hF_1^{\wedge} ; \quad F_1^{\wedge} = F(N) , \quad (5.c)$$

$$q^1 = q^0(x_1) + hF_1^{-} ; \quad F_1^{-} = F(M) .$$

We form the elementary increments:

$$\left. \begin{aligned} \Delta^1 u &= u^1 - u^0(x_1) \\ \Delta^1 p &= p^1 - p^0(x_1) \\ \Delta^1 q &= q^1 - q^0(x_1) \end{aligned} \right\} , \quad (5.d) \quad \left. \begin{aligned} \delta^1 u &= u^1 - u^0(x_2) \\ \delta^1 p &= p^1 - p^0(x_2) \\ \delta^1 q &= q^1 - q^0(x_2) \end{aligned} \right\} . \quad (5.e)$$

Increments (5.d) and (7.d) below relate to the point M, and increments (5.e) — to the point N.

Then, we introduce the intermediate points $M_2^j (j = u, q)$, $N_2^j (j = u, p)$:

$$M_2^j = \left[x_1 + \rho^j h ; y_1 + \sigma^j h ; u^0(x_1) + \alpha^j \Delta^1 u ; \right. \\ \left. p^0(x_1) + \beta^j \Delta^1 p ; \quad q^0(x_1) + \gamma^j \Delta^1 q \right] , \quad (6.a)$$

$$N_2^j = \left[x_2 + \bar{\rho}^j h ; y_2 + \bar{\sigma}^j h ; u^0(x_2) + \bar{\alpha}^j \delta^1 u ; \right. \\ \left. p^0(x_2) + \bar{\beta}^j \delta^1 p ; \quad q^0(x_2) + \bar{\gamma}^j \delta^1 q \right] , \quad (6.b)$$

$\rho^j, \bar{\rho}^j, \sigma^j, \bar{\sigma}^j, \dots, \gamma^j, \bar{\gamma}^j$ are numerical parameters.

At the second iteration we assume that:

$$u^2 = u^0(x_1) + I(h) + \frac{1}{2} h^2 F_2 , \quad (7.a)$$

$$F_2 = \omega_2 F(M_2^u) + \tau_2 F(N_2^u) ; \quad \omega_2, \tau_2 \geq 0 ; \quad \omega_2 + \tau_2 = 1 ,$$

$$p^2 = p^0(x_2) + hF_2^{\wedge} ; \quad F_2^{\wedge} = F(N_2^p) , \quad (7.b)$$

$$q^2 = q^0(x_1) + hF_2^{-} ; \quad F_2^{-} = F(M_2^q) \quad (7.c)$$

$\left[\text{by } F(M_2^j) \right]$ is understood $f[x_1 + \rho^j h; y_1 + \sigma^j h; u^0(x_1) + \alpha^j \Delta^1 u; p^0(x_1) + \beta^j \Delta^1 p; q^0(x_1) + \gamma^j \Delta^1 q]$; analogously with $\left[F(N_2^j) \right]$,

$$\left. \begin{aligned} \Delta^2 u &= u^2 - u^0(x_1) \\ \Delta^2 p &= p^2 - p^0(x_1) \\ \Delta^2 q &= q^2 - q^0(x_1) \end{aligned} \right\} . \quad (7.d)$$

Finally, as Δu , Δp , Δq we take linear combinations:

$$\left. \begin{aligned} \Delta u &= \lambda_1 \Delta^1 u + \lambda_2 \Delta^2 u \\ \Delta p &= \mu_1 \Delta^1 p + \mu_2 \Delta^2 p \\ \Delta q &= \nu_1 \Delta^1 q + \nu_2 \Delta^2 q \end{aligned} \right\} , \quad (8)$$

λ_i, μ_i, ν_i are numerical parameters.

3. Numerical Determination of Runge-Kutta Parameters

For $\Delta^i u$, $\Delta^i p$, $\Delta^i q$ the following expansions in powers of h are valid:

$$\begin{aligned} \Delta^i u &= h p^0(x_1) + \frac{h^2}{2!} \frac{dp^0(x_1)}{dx} + \frac{h^3}{3!} \frac{d^2 p^0(x_1)}{dx^2} + \frac{h^2}{2!} F(M) \\ &\quad \left| + \frac{h^3}{3!} 3 \left(\frac{dF^i}{dh} \right)_M + O(h^4) \right. , \end{aligned} \quad (9.a)$$

$$\begin{aligned} \Delta^i p &= h \frac{dp^0(x_1)}{dx} + \frac{h^2}{2!} \frac{d^2 p^0(x_1)}{dx^2} + h F(M) \\ &\quad + \frac{h^2}{2!} 2 \left(\frac{dF^i}{dh} \right)_M + O(h^3) , \end{aligned} \quad (9.b)$$

$$\Delta^i q = h F(M) + \frac{h^2}{2!} 2 \left(\frac{dF^i}{dh} \right)_M + O(h^3) . \quad (9.c)$$

$(d/dh)_M$ in (9) and below means that the derivative with respect to h is being taken at the point M .

Taking into account (8) we obtain expansions in powers of h for Δu , Δp , Δq :

$$\begin{aligned} \Delta u = & (\lambda_1 + \lambda_2) \left[hp^0(x_1) + \frac{h^2}{2!} \frac{dp^0(x_1)}{dx} + \frac{h^3}{3!} \frac{d^2p^0(x_1)}{dx^2} \right] \\ & + (\lambda_1 + \lambda_2) \frac{h^2}{2!} F(M) + \frac{h^3}{3!} 3 \left(\lambda_1 \frac{dF_1}{dh} + \lambda_2 \frac{dF_2}{dh} \right)_M \\ & + O(h^4) , \end{aligned} \quad (10.a)$$

$$\begin{aligned} \Delta p = & (\mu_1 + \mu_2) h \frac{dp^0(x_1)}{dx} + \frac{h^2}{2!} \frac{d^2p^0(x_1)}{dx^2} + (\mu_1 + \mu_2) hF(M) \\ & + \frac{h^2}{2!} 2 \left(\mu_1 \frac{dF_1^+}{dh} + \mu_2 \frac{dF_2^+}{dh} \right)_M + O(h^3) , \end{aligned} \quad (10.b)$$

$$\Delta q = (\nu_1 + \nu_2) hF(M) + \frac{h^2}{2!} 2 \left(\nu_1 \frac{dF_1^-}{dh} + \nu_2 \frac{dF_2^-}{dh} \right)_M + O(h^3) . \quad (10.c)$$

The selection of parameters is effected so that the expansions (10) and (4) coincide at the arbitrary function $F(x, y)$ and arbitrary step h . As a result of comparison of (10) and (4) we obtain the following conditions:

$$\lambda_1 + \lambda_2 = \mu_1 + \mu_2 = \nu_1 + \nu_2 = 1 , \quad (11)$$

$$3 \left(\lambda_1 \frac{dF_1}{dh} + \lambda_2 \frac{dF_2}{dh} \right)_M = 2 \frac{\partial F(M)}{\partial x} - \frac{\partial F(M)}{\partial y} , \quad (12)$$

$$2 \left(\mu_1 \frac{dF_1^+}{dh} + \mu_2 \frac{dF_2^+}{dh} \right)_M = 2 \frac{\partial F(M)}{\partial x} - \frac{\partial F(M)}{\partial y} , \quad (13)$$

$$2 \left(\nu_1 \frac{dF_1^-}{dh} + \nu_2 \frac{dF_2^-}{dh} \right)_M = \frac{\partial F(M)}{\partial x} . \quad (14)$$

For the right sides of (12) through (14) we have:

$$\frac{\partial F}{\partial x} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial f}{\partial p} \frac{\partial p}{\partial x} + \frac{\partial f}{\partial q} \frac{\partial q}{\partial x} ,$$

$$\frac{\partial F}{\partial y} = \frac{\partial f}{\partial y} + \frac{\partial f}{\partial u} \frac{\partial u}{\partial y} + \frac{\partial f}{\partial p} \frac{\partial p}{\partial y} + \frac{\partial f}{\partial q} \frac{\partial q}{\partial y} ,$$

where

$$\frac{\partial u}{\partial x} = p ; \quad \frac{\partial u}{\partial y} = q ; \quad \frac{\partial p}{\partial y} = \frac{\partial q}{\partial x} = f ;$$

$$\frac{\partial p}{\partial x} = \lim_{h \rightarrow 0} \frac{\Delta^* p}{h} = \frac{dp^0(x)}{dx} + f ;$$

$$\frac{\partial q}{\partial y} = \lim_{h \rightarrow 0} \frac{q^0(x_1) + \Delta^* q - q^0(x_2)}{h} = - \frac{dq^0(x)}{dx} + f . \quad (15)$$

From (15) it follows that:

$$\begin{aligned} 2 \frac{\partial F(M)}{\partial x} - \frac{\partial F(M)}{\partial y} &= \left[2 \frac{\partial f}{\partial x} - \frac{\partial f}{\partial y} + \frac{\partial f}{\partial u} (2p^0 - q^0) \right. \\ &\quad \left. + \frac{\partial f}{\partial p} \left(2 \frac{dp^0}{dx} + f \right) + \frac{\partial f}{\partial q} \left(\frac{dq^0}{dx} + f \right) \right]_M . \end{aligned} \quad (16)$$

For the calculation of the left sides of (12) through (14) we will find $(d/dh)_M$ for $F(N)$, $F(M_2^j)$, $F(N_2^j)$:

$$\frac{dF(N)}{dh} = \frac{dF(x_1 + h, y_1 - h)}{dh} = \frac{\partial F(N)}{\partial x} - \frac{\partial F(N)}{\partial y} ,$$

so that on the basis of (15) we have

$$\left[\frac{d}{dh} F(N) \right]_M = \left[\frac{\partial f}{\partial x} - \frac{\partial f}{\partial y} + \frac{\partial f}{\partial u} (p^0 - q^0) + \frac{\partial f}{\partial p} \frac{dp^0}{dx} + \frac{\partial f}{\partial q} \frac{dq^0}{dx} \right]_M . \quad (17)$$

Furthermore, from (9) it follows that:

$$\left(\frac{d\Delta^1 u}{dh} \right)_M = p^0(x_1) , \quad \left(\frac{d\Delta^1 p}{dh} \right)_M = \frac{dp^0(x_1)}{dx} + f(M) , \quad \left(\frac{d\Delta^1 q}{dh} \right)_M = f(M) , \quad (18)$$

and from comparison of $\delta^1 u$, $\delta^1 p$, $\delta^1 q$ with $\Delta^1 u$, $\Delta^1 p$, $\Delta^1 q$ we obtain:

$$\left(\frac{d\delta^1 u}{dh}\right)_M = q^0(x_1) , \quad \left(\frac{d\delta^1 p}{dh}\right)_M = f(M) , \quad \left(\frac{d\delta^1 q}{dh}\right)_M = -\frac{dq^0(x_1)}{dx} + F(M) . \quad (19)$$

From (6.a), by using (18), we have

$$\begin{aligned} \left[\frac{d}{dh} F(M_2^j)\right]_M &= \left[\rho^j \frac{\partial f}{\partial x} + \sigma^j \frac{\partial f}{\partial y} + \alpha^j p^0 \frac{\partial f}{\partial u} \right. \\ &\quad \left. + \beta^j \left(\frac{dp^0}{dx} + f\right) \frac{\partial f}{\partial p} + \gamma^j f \frac{\partial f}{\partial q}\right]_M . \end{aligned} \quad (20)$$

Analogously, from (6.b), where $x_2 = x_1 + h$, $y_2 = y_1 - h$, by using (19), we have

$$\begin{aligned} \left[\frac{d}{dh} F(N_2^j)\right]_M &= \left[\left(1 + \bar{\rho}^j\right) \frac{\partial f}{\partial x} + \left(\bar{\sigma}^j - 1\right) \frac{\partial f}{\partial y} + \left(p^0 - q^0 + \bar{\alpha}^j q^0\right) \frac{\partial f}{\partial u} \right. \\ &\quad \left. + \left(\frac{dp^0}{dx} + \beta^j f\right) \frac{\partial f}{\partial p} + \left(\frac{dq^0}{dx} - \gamma^j \frac{dq^0}{dx} + \bar{\gamma}^j f\right) \frac{\partial f}{\partial q}\right]_M . \end{aligned} \quad (21)$$

Let us return to the expression (12). The right side of this equality is given in (16).

By using (5.a), (7.a), (17), (20), and (21), its left side can be expressed in the following form:

$$\begin{aligned} &3 \frac{\partial f}{\partial x} (\lambda_1 \tau_1 + \lambda_2 \tau_2 + \lambda_2 \omega_2 \rho + \lambda_2 \tau_2 \bar{\rho}) + 3 \frac{\partial f}{\partial y} (-\lambda_1 \tau_1 - \lambda_2 \tau_2 + \lambda_2 \omega_2 \sigma + \lambda_2 \tau_2 \bar{\sigma}) \\ &+ 3 \frac{\partial f}{\partial u} [(\lambda_1 \tau_1 + \lambda_2 \tau_2 + \lambda_2 \omega_2 \alpha) p^0 - (\lambda_1 \tau_1 + \lambda_2 \tau_2 - \lambda_2 \tau_2 \bar{\alpha}) q^0] \\ &+ 3 \frac{\partial f}{\partial p} \left[(\lambda_1 \tau_1 + \lambda_2 \tau_2 + \lambda_2 \omega_2 \beta) \frac{dp^0}{dx} + (\lambda_2 \omega_2 \beta + \lambda_2 \tau_2 \bar{\beta}) f \right] \\ &+ 3 \frac{\partial f}{\partial q} \left[(\lambda_1 \tau_1 + \lambda_2 \tau_2 - \lambda_2 \tau_2 \bar{\gamma}) \frac{dq^0}{dx} + (\lambda_2 \omega_2 \gamma + \lambda_2 \tau_2 \bar{\gamma}) f \right]. \end{aligned} \quad (22)$$

Here all values are given at the point M and, for simplicity, all indices $j = u$ are omitted. Assume:

$$\bar{\alpha} = \alpha , \quad \bar{\beta} = \beta , \quad \bar{\gamma} = \gamma , \quad \bar{\rho} = \rho - 1 , \quad \bar{\sigma} = \sigma + 1 . \quad (23)$$

(Such a choice of $\bar{\sigma}$, $\bar{\rho}$ is dictated by the form of the region MNP; with this choice the corresponding projections of points M_2^u and N_2^u in the plane xy coincide.)

Taking into account (11), (23) and equating (22) to the right side of (16) we obtain the following system of equations for the parameters at Δu :

$$\lambda_1 \tau_1 + \lambda_2 \rho = \frac{2}{3} , \quad \lambda_1 \tau_1 - \lambda_2 \sigma = \frac{1}{3} ,$$


$$\lambda_1 \tau_1 + \lambda_2 \tau_2 + \lambda_2 \omega_2 \alpha = \lambda_1 \tau_1 + \lambda_2 \tau_2 + \lambda_2 \omega_2 \beta = \frac{2}{3} ,$$

$$\lambda_1 \tau_1 + \lambda_2 \tau_2 - \lambda_2 \tau_2 \alpha = \lambda_1 \tau_1 + \lambda_2 \tau_2 - \lambda_2 \tau_2 \gamma = \frac{1}{3} ,$$

$$\lambda_2 \beta = \lambda_2 \gamma = \frac{1}{3} . \quad (24)$$

For example, the parameters from Table I will constitute solutions of the system (24).

TABLE I

	ω_1	τ_1	ω_2	τ_2	λ_1	λ_2	ρ	σ	$\bar{\rho}$	$\bar{\sigma}$	$\alpha, \bar{\alpha}$	$\beta, \bar{\beta}$	$\gamma, \bar{\gamma}$	Location of Projections of M_2^u and N_2^u in the Plane xy
1)	0	1	1	0	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{1}{2}$	0	-	-	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	
2)	1	0	0	1	$\frac{1}{3}$	$\frac{2}{3}$	-	-	0	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	
3)	1	0	$\frac{1}{2}$	$\frac{1}{2}$	0	1	$\frac{2}{3}$	$-\left(\frac{1}{3}\right)$	$-\left(\frac{1}{3}\right)$	$\frac{2}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	
4)	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{5}{6}$	$-\left(\frac{1}{6}\right)$	$-\left(\frac{1}{6}\right)$	$\frac{5}{6}$	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	

Analogously, at Δp and Δq we obtain for the parameters:

$$\mu_2 \bar{\rho}^p = 0, \quad \mu_2 \bar{\sigma}^p = \frac{1}{2}, \quad \mu_2 \bar{\alpha}^p = \mu_2 \bar{\beta}^p = \mu_2 \bar{\gamma}^p = \frac{1}{2}, \quad (25)$$

$$\nu_2 \rho^q = \frac{1}{2}, \quad \sigma^q = 0, \quad \nu_2 \alpha^q = \nu_2 \beta^q = \nu_2 \gamma^q = \frac{1}{2}. \quad (26)$$

Equations (25) and (26) can be satisfied upon selection of parameters, for example, from variants in Table II.

TABLE II

μ_1, ν_1	μ_2, ν_2	$\bar{\rho}^p, \sigma^q$	$\bar{\sigma}^p, \rho^q$	$\bar{\alpha}^p, \alpha^q$	$\bar{\beta}^p, \beta^q$	$\bar{\gamma}^p, \gamma^q$
0	1	0	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$\frac{1}{2}$	$\frac{1}{2}$	0	1	1	1	1

The factual scheme of calculations according to proposed algorithms has the form:

$$\left. \begin{aligned} u(P) &= u^0(x_1) + \lambda_1 \Delta^1 u + \lambda_2 \Delta^2 u, \\ p(P) &= p^0(x_2) + \mu_1 \delta^1 p + \mu_2 \delta^2 p, \\ q(P) &= q^0(x_1) + \nu_1 \Delta^1 q + \nu_2 \Delta^2 q, \end{aligned} \right\}. \quad (27)$$

where

$$\left. \begin{aligned} \Delta^1 u &= I(h) + \frac{1}{2} h^2 F_1, \\ \delta^1 p &= h F_1^+, \\ \Delta^1 q &= h F_1^-; \end{aligned} \right\} \quad (28.a) \quad \left. \begin{aligned} \delta^1 u &= \Delta^1 u + u^0(x_1) - u^0(x_2), \\ \Delta^1 p &= \delta^1 p + p^0(x_2) - p^0(x_1), \\ \delta^1 q &= \Delta^1 q + q^0(x_1) - q^0(x_2); \end{aligned} \right\} \quad (28.b)$$

$$\left. \begin{aligned} \Delta^2 u &= I(h) + \frac{1}{2} h^2 F_2 , \\ \delta^2 p &= h F_2^+ , \\ \Delta^2 q &= h F_2^- . \end{aligned} \right\} \quad (28.c)$$

If we adopt the first variants of Tables I and II, then at each step we will have to calculate only four values of f : $f(M)$, $f(N)$, $f(M_2^u) = f(M_2^q)$, $f(N_2^p)$.

4. Investigation of the Stability of the Method

We find $\int_{x_1}^{x_1 + h} p^0(x) dx = I(h)$ according to Simpson's formula

$$I(h) \approx \frac{1}{6} h [p^0(M) + 4p^0(Q) + p^0(N)] ,$$

where Q is the midpoint of the segment MN . We calculate the unknown $p^0(Q)$ with the aid of Bessel's interpolation polynomial or, if need be, with Newton's polynomial. Whereupon, if the values of $p^0(x)$ in the nodes of the given layer are found with an error η , and if we neglect the remainder of order $O(h^5)$ in Simpson's formula, then the irremovable error in the calculation of $I(h)$ will be approximately $h\eta$.

Relationships (27) on the k^{th} layer will have the following form:

$$u_k = u_{k-1} + \lambda_1 \Delta^1 u_{k-1} + \lambda_2 \Delta^2 u_{k-1} , \quad (29.a)$$

$$p_k = p_{k-1} + \mu_1 \delta^1 p_{k-1} + \mu_2 \delta^2 p_{k-1} , \quad (29.b)$$

$$q_k = q_{k-1} + \nu_1 \Delta^1 q_{k-1} + \nu_2 \Delta^2 q_{k-1} , \quad (29.c)$$

where the indices k relate to the point P of the triangle MNP , indices $k-1$ in (29.a), (29.c) relate to the point M , and in (29.b) relate to the point N .

Along with (29) we write:

$$u_k^* = u_{k-1}^* + \lambda_1 \Delta^1 u_{k-1}^* + \lambda_2 \Delta^2 u_{k-1}^* + r_1 , \quad (30.a)$$

$$p_k^* = p_{k-1}^* + \mu_1 \delta^1 p_{k-1}^* + \mu_2 \delta^2 p_{k-1}^* + r_2, \quad (30.b)$$

$$q_k^* = q_{k-1}^* + \nu_1 \Delta^1 q_{k-1}^* + \nu_2 \Delta^2 q_{k-1}^* + r_3. \quad (30.c)$$

Here, $\Delta^1 u_{k-1}^*$, $\Delta^2 u_{k-1}^*$, $\delta^1 p_{k-1}^*$, ..., $\Delta^2 q_{k-1}^*$ are elementary increments calculated with respect to (28.a) through (28.c), in which the approximate values u , p , q are replaced by the exact values u^* , p^* , q^* ; r_1 , r_2 , r_3 are the local errors of the method which in our case are of orders $O(h^4)$, $O(h^3)$, $O(h^3)$ [equation (10)].

We denote the errors as follows:

$$|u_k^* - u_k| = \epsilon_k, \quad |p_k^* - p_k| = \eta_k, \quad |q_k^* - q_k| = \Theta_k. \quad (31)$$

Subtracting (29) from (30) and using designations (31) we obtain:

$$\left. \begin{aligned} \epsilon_k &\leq \epsilon_{k-1} + \lambda_1 |\Delta^1 u_{k-1}^* - \Delta^1 u_{k-1}| + \lambda_2 |\Delta^2 u_{k-1}^* - \Delta^2 u_{k-1}| + r_1, \\ \eta_k &\leq \eta_{k-1} + \mu_1 |\delta^1 p_{k-1}^* - \delta^1 p_{k-1}| + \mu_2 |\delta^2 p_{k-1}^* - \delta^2 p_{k-1}| + r_2, \\ \Theta_k &\leq \Theta_{k-1} + \nu_1 |\Delta^1 q_{k-1}^* - \Delta^1 q_{k-1}| + \nu_2 |\Delta^2 q_{k-1}^* - \Delta^2 q_{k-1}| + r_3, \end{aligned} \right\} \quad (32)$$

Let $f(x, y, u, p, q)$ satisfy a Lipschitz condition with respect to u , p , q , with constant K , and let

$$U = K(\epsilon_{k-1} + \eta_{k-1} + \Theta_{k-1}).$$

On the strength of (28.a) we have

$$\left. \begin{aligned} |\Delta^1 u_{k-1}^* - \Delta^1 u_{k-1}| &\leq h \eta_{k-1} + \frac{1}{2} h^2 U, \\ |\delta^1 p_{k-1}^* - \delta^1 p_{k-1}| &\leq h U, \quad |\Delta^1 q_{k-1}^* - \Delta^1 q_{k-1}| \leq h U. \end{aligned} \right\} \quad (33)$$

Furthermore, from (33) and (28.b) it follows that:

$$|\delta^1 u_{k-1}^* - \delta^1 u_{k-1}| \leq 2\epsilon_{k-1} + h\eta_{k-1} + \frac{1}{2}h^2U ,$$

$$|\Delta^1 p_{k-1}^* - \Delta^1 p_{k-1}| \leq 2\eta_{k-1} + hU ,$$

$$|\delta^1 q_{k-1}^* - \delta^1 q_{k-1}| \leq 2\Theta_{k-1} + hU .$$

Taking into account (6.a) and (6.b) we have:

$$|f(M_2^{j*}) - f(M_2^j)| \leq V = K \left[\left(\epsilon_{k-1} + h\eta_{k-1} + \frac{1}{2}h^2U \right) + \left(3\eta_{k-1} + hU \right) + \left(\Theta_{k-1} + hU \right) \right] ;$$

$$|f(N_2^{j*}) - f(N_2^j)| \leq W = K \left[\left(3\epsilon_{k-1} + h\eta_{k-1} + \frac{1}{2}h^2U \right) + \left(\eta_{k-1} + hU \right) + \left(3\Theta_{k-1} + hU \right) \right] .$$

From the preceding and from the equality (28,c) we obtain:

$$\left. \begin{aligned} |\Delta^2 u_{k-1}^* - \Delta^2 u_{k-1}| &\leq h\eta_{k-1} + \frac{1}{2}h^2(V + W) , \\ |\delta^2 p_{k-1}^* - \delta^2 p_{k-1}| &\leq hW , \quad |\Delta^2 q_{k-1}^* - \Delta^2 q_{k-1}| \leq hV . \end{aligned} \right\} \quad (34)$$

Using (33) and (34) we write (32) in the form:

$$\begin{aligned} \epsilon_k &\leq \epsilon_{k-1} + h\eta_{k-1} + \frac{1}{2}h^2[\lambda_1 U + \lambda_2(V + W)] + r_1 , \\ \eta_k &\leq \eta_{k-1} + h(\mu_1 U + \mu_2 W) + r_2 , \\ \Theta_k &\leq \Theta_{k-1} + h(\nu_1 U + \nu_2 V) + r_3 . \end{aligned} \quad (35)$$

We denote vectors with components ϵ_k , η_k , Θ_k and r_1 , r_2 , r_3 correspondingly by X_k and r ($r \sim \{O(h^4), O(h^3), O(h^3)\}$), and we assume that they are accurately given on the zero layer of u^0 , p^0 , q^0 . On the basis of (35) we have

$$X_k \leq AX_{k-1} + r, \quad k = 1, 2, \dots, \left\lceil \frac{AC}{h} \right\rceil.$$

Diagonal elements of matrices of A are

$$1 + c_1 h^2 + \dots, \quad 1 + c_2 h + \dots, \quad 1 + c_3 h + \dots,$$

where c_1, c_2, c_3 are constants not depending on h ; the dots designate certain addends of higher powers in h ; non-diagonal elements of A are values of the order not higher than $O(h)$.

Thus, at usual norms of vectors and matrices the distribution coefficient of errors at each step does not exceed $1 + O(h)$; hence ensues the stability of the method in the sense of [4].

5. Some Generalizations of the Method

Let there be a system of n equations

$$u_{xy} = f(x, y, u, p, q) \quad (36)$$

with initial conditions:

$$u^0 = u^0(x), \quad p^0 = p^0(x), \quad q^0 = q^0(x), \quad (37)$$

given along the segment AB of the straight line $x + y = \text{const}$, where u is n -dimensional vector function of x, y ; $p = u_x$; $q = u_y$. All that was stated in

Paragraphs 2 and 3 remains valid for the problem (36) and (37), with the difference that all functions and increments in algorithms (27), (28) are

vectorial-values depending on the scalar step h ; $\lambda_i, \mu_i, \nu_i, \alpha^i, \beta^i, \gamma^i, \bar{\alpha}^i, \bar{\beta}^i, \bar{\gamma}^i, \rho^j, \bar{\rho}^j, \sigma^j, \bar{\sigma}^j$ are scalar coefficients. The proof is analogous to that carried out in Paragraphs 2 and 3; there remain in force conditions (11), (24 through (26) and the results of Tables I and II.

Let us apply the stated method to the solution of the Cauchy problem for nonlinear equations of the hyperbolic type with two independent variables by reducing such equations to the system (36) with initial conditions (37). Let us consider, for example, the Cauchy problem for the Monge-Ampere equation

$$ar + 2bs + ct + g(rt - s^2) + f = 0, \quad (g \neq 0), \quad (38)$$

where $r = u_{xx}$; $s = u_{xy}$; $t = u_{yy}$; u is an unknown function of x, y ; a, b, c, g, f are given functions of x, y, u, p, q , while $p = u_x$ and $q = u_y$. Assume that initial conditions are given along the segment A'B' of a noncharacteristic curve Γ in the form of $u^0 = u^0(x)$, $p^0 = p^0(x)$, $q^0 = q^0(x)$.

It is known [5] that in the hyperbolic case the equation (38) is equivalent to the system

$$\left. \begin{aligned} c \frac{\partial x}{\partial \alpha} + \varphi_1 \frac{\partial y}{\partial \alpha} + g \frac{\partial p}{\partial \alpha} &= 0, \\ \varphi_2 \frac{\partial x}{\partial \alpha} + \alpha \frac{\partial y}{\partial \alpha} + g \frac{\partial q}{\partial \alpha} &= 0, \\ p \frac{\partial x}{\partial \alpha} + q \frac{\partial y}{\partial \alpha} - \frac{\partial u}{\partial \alpha} &= 0, \\ c \frac{\partial x}{\partial \beta} + \varphi_2 \frac{\partial y}{\partial \beta} + g \frac{\partial p}{\partial \beta} &= 0, \\ \varphi_1 \frac{\partial x}{\partial \beta} + \alpha \frac{\partial y}{\partial \beta} + g \frac{\partial q}{\partial \beta} &= 0, \end{aligned} \right\} \quad (39)$$

where α, β are parameters of characteristic families; φ_1, φ_2 are roots of the characteristic equation $\varphi^2 - 2b\varphi + ac - gf = 0$, whereas φ_1 corresponds to a family α , and φ_2 to a family β .

We will differentiate the first three equations of the system (39) in accordance with Levi and Friedrichs, with respect to β , and the remainder of them — with respect to α , and we will solve the system thus obtained in terms of the second derivatives. As a result we will obtain a system composed of five equations:

$$x_{\alpha\beta} = f_1, \quad y_{\alpha\beta} = f_2, \quad u_{\alpha\beta} = f_3, \quad p_{\alpha\beta} = f_4, \quad q_{\alpha\beta} = f_5, \quad (40)$$

where f_j ($j = 1 \div 5$) are known functions of $x, y, u, p, q, x_\alpha, y_\alpha, u_\alpha, p_\alpha, q_\alpha, x_\beta, y_\beta, u_\beta, p_\beta, q_\beta$.

We will select the transformation $(x, y) \rightarrow (\alpha, \beta)$ so that the segment A'B' of the initial curve is transformed into the segment AB of the straight line $\alpha + \beta = \text{const}$. Along AB we know x^0, y^0, u^0, p^0, q^0 and we can uniquely

determine $x_\alpha^0, y_\alpha^0, u_\alpha^0, p_\alpha^0, q_\alpha^0, x_\beta^0, y_\beta^0, \dots, q_\beta^0$. (Details of the stated facts have been given previously [6, 7]). Thus, the Cauchy problem for (38) was reduced to the problem (36), (37); hence follows its solvability by means of the proposed algorithms.

BIBLIOGRAPHY

1. J. Albrecht, ZAMM, 37, 1957, p. 202.
2. G. Strang, NUMERISCHE MATHEMATIK, 6, No. 1, 1964, p. 37.
3. H. Moore, ARCHIVES RAT. MECH. ANALYSIS, 7, 1961, p. 37.
4. V. S. Ryabenkiy and A. F. Filippov, ON THE STABILITY OF DIFFERENCE EQUATIONS (Ob ustoychivosti raznostnykh uravneniy), Moscow, 1956.
5. R. Sauer, ANFANGSWERTPROBLEME BEI PARTIELLEN DIFFERENTIALGLEICHUNGEN, Berlin, 1958.
6. R. Kurant and K. Friedrichs, THE SUPERSONIC FLOW AND SHOCK WAVES (Sverkhzvukovoye techeniye i udarnyye volny), IL, 1950.
7. V. Vazov and D. Forsayt, DIFFERENCE METHODS FOR SOLVING PARTIAL DIFFERENTIAL EQUATIONS (Razrostnyye metody resheniya differentsial'nykh uravneniy v chastnykh proizvodnykh), IL, 1963.

DISTRIBUTION

	No. of Copies		No. of Copies
<u>EXTERNAL</u>		U. S. Atomic Energy Commission	1
Air University Library	1	ATTN: Reports Library, Room G-017	
ATTN: AUL3T		Washington, D. C. 20545	
Maxwell Air Force Base, Alabama 36112		U. S. Naval Research Laboratory	1
U. S. Army Electronics Proving Ground	1	ATTN: Code 2027	
ATTN: Technical Library		Washington, D. C. 20390	
Fort Huachuca, Arizona 85613		Weapons Systems Evaluation Group	1
Naval Weapons Center	1	Washington, D. C. 20305	
ATTN: Technical Library, Code 753		John F. Kennedy Space Center, NASA	2
China Lake, California 93555		ATTN: KSC Library, Documents Section	
Naval Weapons Center, Corona Laboratories	1	Kennedy Space Center, Florida 32899	
ATTN: Documents Librarian		APGC (PGBPS-12)	1
Corona, California 91720		Eglin Air Force Base, Florida 32542	
Lawrence Radiation Laboratory	1	U. S. Army CDC Infantry Agency	1
ATTN: Technical Information Division		Fort Benning, Georgia 31905	
P. O. Box 808		Argonne National Laboratory	1
Livermore, California 94550		ATTN: Report Section	
Sandia Corporation	1	9700 South Cass Avenue	
ATTN: Technical Library		Argonne, Illinois 60440	
P. O. Box 969		U. S. Army Weapons Command	1
Livermore, California 94551		ATTN: AMSWE-RDR	
U. S. Naval Postgraduate School	1	Rock Island, Illinois 61201	
ATTN: Library		Rock Island Arsenal	1
Monterey, California 93940		ATTN: SWERI-RDI	
Electronic Warfare Laboratory, USAECOM	1	Rock Island, Illinois 61201	
Post Office Box 205		U. S. Army Cmd. & General Staff College	1
Mountain View, California 94042		ATTN: Acquisitions, Library Division	
Jet Propulsion Laboratory	2	Fort Leavenworth, Kansas 66027	
ATTN: Library (TDS)		Combined Arms Group, USACDC	1
4800 Oak Grove Drive		ATTN: Op. Res., P and P Div.	
Pasadena, California 91103		Fort Leavenworth, Kansas 66027	
U. S. Naval Missile Center	1	U. S. Army CDC Armor Agency	1
ATTN: Technical Library, Code N3022		Fort Knox, Kentucky 40121	
Point Mugu, California 93041		Michoud Assembly Facility, NASA	1
U. S. Army Air Defense Command	1	ATTN: Library, I-MICH-OSD	
ATTN: ADSX		P. O. Box 29300	
Ent Air Force Base, Colorado 80912		New Orleans, Louisiana 70129	
Central Intelligence Agency	4	Aberdeen Proving Ground	1
ATTN: OCR/DD-Standard Distribution		ATTN: Technical Library, Bldg. 313	
Washington, D. C. 20505		Aberdeen Proving Ground, Maryland 21005	
Harry Diamond Laboratories	1	NASA Sci. & Tech. Information Facility	5
ATTN: Library		ATTN: Acquisitions Branch (S-AK/DL)	
Washington, D. C. 20438		P. O. Box 33	
Scientific & Tech. Information Div., NASA	1	College Park, Maryland 20740	
ATTN: ATS		U. S. Army Edgewood Arsenal	1
Washington, D. C. 20546		ATTN: Librarian, Tech. Info. Div.	
		Edgewood Arsenal, Maryland 21010	

	No. of Copies		No. of Copies
National Security Agency ATTN: C3/TDL Fort Meade, Maryland 20755	1	Brookhaven National Laboratory Technical Information Division ATTN: Classified Documents Group Upton, Long Island, New York 11973	1
Goddard Space Flight Center, NASA ATTN: Library, Documents Section Greenbelt, Maryland 20771	1	Watervliet Arsenal ATTN: SWEWV-RD Watervliet, New York 12189	1
U. S. Naval Propellant Plant ATTN: Technical Library Indian Head, Maryland 20640	1	U. S. Army Research Office (ARO-D) ATTN: CRD-AA-IP Box CM, Duke Station Durham, North Carolina 27706	1
U. S. Naval Ordnance Laboratory ATTN: Librarian, Eva Liberman Silver Spring, Maryland 20910	1	Lewis Research Center, NASA ATTN: Library 21000 Brookpark Road Cleveland, Ohio 44135	1
Air Force Cambridge Research Labs. L. G. Hanscom Field ATTN: CRMXL/Stop 29 Bedford, Massachusetts 01730	1	Foreign Technology Division ATTN: Library Wright-Patterson Air Force Base, Ohio 45400	1
U. S. Army Tank Automotive Center ATTN: SMOTA-RTS.1 Warren, Michigan 48090	1	U. S. Army Artillery & Missile School ATTN: Guided Missile Department Fort Sill, Oklahoma 73503	1
U. S. Army Materials Research Agency ATTN: AMXMR-ATL Watertown, Massachusetts 02172	1	U. S. Army CDC Artillery Agency ATTN: Library Fort Sill, Oklahoma 73504	1
Strategic Air Command (OAI) Offutt Air Force Base, Nebraska 68113	1	U. S. Army War College ATTN: Library Carlisle Barracks, Pennsylvania 17013	1
Picatinny Arsenal, USAMUCOM ATTN: SMUPA-VA6 Dover, New Jersey 07801	1	U. S. Naval Air Development Center ATTN: Technical Library Johnsville, Warminster, Pennsylvania 18974	1
U. S. Army Electronics Command ATTN: AMSEL-CB Fort Monmouth, New Jersey 07703	1	Frankford Arsenal ATTN: C-2500-Library Philadelphia, Pennsylvania 19137	1
Sandia Corporation ATTN: Technical Library P. O. Box 5800 Albuquerque, New Mexico 87115	1	Div. of Technical Information Ext., USAEC P. O. Box 62 Oak Ridge, Tennessee 37830	1
ORA(RRRT) Holloman Air Force Base, New Mexico 88330	1	Oak Ridge National Laboratory ATTN: Central Files P. O. Box X Oak Ridge, Tennessee 37830	1
Los Alamos Scientific Laboratory ATTN: Report Library P. O. Box 1663 Los Alamos, New Mexico 87544	1	Air Defense Agency, USACDC ATTN: Library Fort Bliss, Texas 79916	1
White Sands Missile Range ATTN: Technical Library White Sands, New Mexico 88002	1	U. S. Army Air Defense School ATTN: AKBAAS-DR-R Fort Bliss, Texas 79906	1
Rome Air Development Center (EMLAL-1) ATTN: Documents Library Griffiss Air Force Base, New York 13440	1		

	No. of Copies		No. of Copies
U. S. Army Combat Developments Command Institute of Nuclear Studies Fort Bliss, Texas 79916	1	<u>INTERNAL</u>	
Manned Spacecraft Center, NASA ATTN: Technical Library, Code BM6 Houston, Texas 77058	1	Headquarters U. S. Army Missile Command Redstone Arsenal, Alabama 35809 ATTN: AMSMI-D	1
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	20	AMSMI-XE, Mr. Lowers	1
U. S. Army Research Office ATTN: STINFO Division 3045 Columbia Pike Arlington, Virginia 22204	1	AMSMI-Y	1
		AMSMI-R, Mr. McDaniel	1
		AMSMI-RAP	1
		AMSMI-RBLD	10
		USACDC-LnO	1
		AMSMI-RB, Mr. Croxton	1
		AMSMI-RBT	8
U. S. Naval Weapons Laboratory ATTN: Technical Library Dahlgren, Virginia 22448	1	National Aeronautics & Space Administration Marshall Space Flight Center Marshall Space Flight Center, Ala. 35812	
U. S. Army Engineer Res. & Dev. Labs. ATTN: Scientific & Technical Info. Br. Fort Belvoir, Virginia 22060	2	ATTN: MS-T, Mr. Wiggins	5
		R-COMP-T, Dr. Fehlberg	1
Langley Research Center, NASA ATTN: Library, MS-185 Hampton, Virginia 23365	1		
Research Analysis Corporation ATTN: Library McLean, Virginia 22101	1		
Foreign Science & Technology Center Munitions Building Washington, D. C. 20315	3		
National Aeronautics & Space Administration Code USS-T (Translation Section) Washington, D. C. 20546	2		

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Redstone Scientific Information Center Research and Development Directorate U. S. Army Missile Command Redstone Arsenal, Alabama 35809		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP N/A	
3. REPORT TITLE NUMERICAL SOLUTION OF HYPERBOLIC EQUATIONS AND SYSTEMS WITH TWO INDEPENDENT VARIABLES BY A METHOD OF THE RUNGE-KUTTA TYPE. I: Vestsi Akademii Navuk Belaruskay SSR, Seryya Fizika Matematychnykh Navuk, No.4, pp. 110-120, 1965			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Translated from the Russian			
5. AUTHOR(S) (First name, middle initial, last name) Nguyen Kong Tuy			
6. REPORT DATE 24 June 1968		7a. TOTAL NO. OF PAGES 22	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. N/A		9a. ORIGINATOR'S REPORT NUMBER(S) RSIC-811	
b. PROJECT NO. N/A		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) AD _____	
c. N/A			
d.			
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES None		12. SPONSORING MILITARY ACTIVITY Same as No. 1	
13. ABSTRACT Discussed is the application of two-iteration Runge-Kutta algorithms in the solution of the Cauchy problem for hyperbolic equations and systems with two independent variables.			

DD FORM 1473
1 NOV 65REPLACES DD FORM 1473, 1 JAN 64, WHICH IS
OBSOLETE FOR ARMY USE.UNCLASSIFIED
Security Classification

21

UNCLASSIFIED

Security Classification

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Runge-Kutta algorithm Cauchy problem Hyperbolic equations Simpson's formula Monge-Ampere equation						

UNCLASSIFIED

Security Classification